

BINOMIAL THEOREM AND EXPONENT FOR VARIABLES COMMUTING AS $yx = qxy$

A. V. STOYANOVSKY

ABSTRACT. We state analogs of the binomial theorem and the exponential function for variables x, y commuting as $yx = qxy$.

Let x, y be formal variables satisfying the commutation relation $yx = qxy$, where q is a number. In this note we are going to write out the formulas for $(x + y)^n$ and for the function (formal series) $\exp_q(x)$ satisfying the functional equation of the exponent. Note that the usual approach to q -analysis is somewhat another (see, for example, [1,2]).

The author is grateful to V. V. Dolotin and A. N. Zhukov for stimulating questions. Thanks are also due to Professors Tom Koornwinder and Uwe Franz, who informed me that this result is not new, but has been discovered by Schutzenberger in 1953 [3], see also [4,5] for further information and references.

Theorem 1.

$$(1) \quad (x + y)^n = \sum_{k=0}^n \binom{n}{k}_q x^k y^{n-k},$$

where $\binom{n}{k}_q$ are the usual q -binomial coefficients (Gaussian polynomials),

$$(2) \quad \binom{n}{k}_q = \frac{(n!)_q}{(k!)_q ((n-k)!)_q},$$

where

$$(3) \quad (n!)_q = (1)_q (2)_q (3)_q \dots (n)_q, \quad (k)_q = 1 + q + q^2 + \dots + q^{k-1},$$

satisfying the recurrence relation

$$(4) \quad \binom{n}{k}_q = \binom{n-1}{k}_q + q^{n-k} \binom{n-1}{k-1}_q, \quad \binom{n}{0}_q = \binom{n}{n}_q = 1,$$

and the symmetry relation

$$(5) \quad \binom{n}{k}_q = \binom{n}{n-k}_q.$$

Theorem 2. *If we put*

$$(6) \quad \exp_q(x) = \sum_{n=0}^{\infty} \frac{x^n}{(n!)_q},$$

then

$$(7) \quad \exp_q(x + y) = \exp_q(x) \exp_q(y).$$

Proof of Theorem 1 is by induction. Theorem 2 follows from Theorem 1.

REFERENCES

- [1] G. Gasper, M. Rahman, Basic hypergeometric series, Encyclopedia of Mathematics and Its Applications, vol. 35, Cambridge University Press, Cambridge, 1990.
- [2] G. E. Andrews, The theory of partitions, Encyclopedia of Mathematics and Its Applications, vol. 2, Addison-Wesley, Reading, Mass.; reissued by Cambridge University Press, Cambridge, 1985.
- [3] M. P. Schutzenberger, Une interprétation de certaines solutions de l'équation fonctionnelle $F(x+y) = F(x)F(y)$, C. R. Acad. Sci. Paris, 236, 352–353 (1953).
- [4] P. Feinsilver, Discrete analogues of the Heisenberg–Weyl algebra, Monatsh. Math. 104, 89–108 (1987).
- [5] T. H. Koornwinder, Special functions and q -commuting variables, arxiv: q-alg/9608008, in: Special Functions, q -Series and Related Topics, M. E. H. Ismail, D. R. Masson and M. Rahman eds., Fields Institute Communications 14, American Mathematical Society, 1997, pp. 131–166.

RUSSIAN STATE UNIVERSITY OF HUMANITIES

E-mail address: stoyan@mccme.ru